



EADU TANTA FARMS AND TRAINING CENTER

AGE IN DAYS	AGE IN WEEKS	VACCINE	DRUGS AND REMARKS
1 st	1 ST WEEK		Glucose
2 nd – 6 th			Antibiotic Vitamins
7 th		1ST GUMBORO(Intermediate)	Vitamins
8 th -10 th	2 ND WEEK		Vitamins
11 th -13 th			Coccidiostat
14 th		1st NEW CASTLE (HB1)	Vitamins
15 th -16 th	3 RD WEEK		Vitamins
17 th			Plain water
18 th – 20 th			Coccidiostat
21 st		2nd GUMBORO (intermediate plus)	Anti-stress
22 nd	4 TH WEEK		Plain water
23 th -25 th			Coccidiostat
26 th			Plain water

27 th			Deworming
28 th		2nd NEW CASTLE (LASOTA)	Anti-stress
29 TH	5 TH WEEK		Plain water
30 TH – 32 nd			Coccidiostat
33 RD -34 th			Vitamins
35 TH		2nd Gumboro (Intermediate plus)	Vitamins
36 th – 38 th	6 TH WEEK		Coccidiostat
39 th – 42 nd			Antibiotic Vitamins
43 th – 45 th	7 TH WEEK		Coccidiostat
46 th -48 th			Vitamins
49 th			Plain water
50 th – 52 nd	8 TH WEEK		Coccidiostat
53 rd -55 th			Vitamins
56 th		1ST FOWL POX	Anti-stress
57 th	9 TH WEEK		Plain water
58 th – 60 th			Coccidiostat
61 st			Deworming
62 nd – 63 rd			Plain water
64 th – 66 th	10 th WEEK		Coccidiostat
67 th – 69 th			Plain water
70 th		2nd LASOTA	Vitamins
71 st	11 th WEEK		Plain water
72 nd -74 th			Coccidiostat
75 th -77 th			Vitamins

78 TH -80 th	12 TH WEEK		Coccidiostat
81 st - 83 rd			Antibiotic Vitamin
84 TH		2ND FOWL POX /DEBEAKING	Anti-stress
85 TH	13 TH WEEK		Plain water
86 th – 88 th			Coccidiostat
89 th - 91 ST			Vitamins
92 nd – 94 th	14 TH WEEK		Coccidiostat
95 th -98 th			Pure Vitamin
99 TH -101 st	15 TH WEEK		Coccidiostat
102 nd -104 th			Vitamins
105 TH			Plain water
106 TH	16 TH WEEK		Plain water
107 th -109 th			Coccidiostat
110 th -111 th			Vitamins
112 th		3RD Newcastle NB: maybe given earlier if Newcastle is endemic in the area	Anti-stress

PLEASE NOTE:

1. This program acts as a guide and can be adjusted as and when circumstances necessitate such changes or additions, with the help of your veterinary advisor.

2. Ensure you have at least 2 types of coccidiostat, Amprolium based and Sulphur based. Do well to alternate between these two coccidiostat
3. Debeaking at 8 or 15 weeks is recommended but it can also be done on the 4th day when the equipment is available. However, where pecking gets rampant, Debeaking can be arranged to reduce the incident.
4. The use of antibiotics should be very specific following a Vets prescription and possibly backed by sensitivity test from the vet lab.
5. Regular cleaning and disinfection of feeders, drinkers and the pen is strongly recommended.
6. Deworm regularly every TWO months and give anti-stress after each deworming.
7. Check for lice, fleas, mites and ticks occasionally, and delouse when necessary. Give anti-stress after treatment.
8. Seek veterinary professional advice and don't give drug haphazardly to your birds.
9. Give anti-stress after vaccination.
10. After 3rd NCD vaccine, repeat after every THREE months.

Antibiotic	Vitamins	Coccidiostat	Dewormer
Tetracolivit	Supervitamin	Amprolium 20S	Levasol
Keproceryl	Aminovit	Amprolium 250	Piper Dewormer
Penprovit	Amintotal Powervit	Franzuril	Franmec Oral
Penstrep powder	Vitamino Trace Oral	Keptox	Keptomec Oral
Neoxyvital Forte	Vitamin B- Choline	Cox B3	
Franceryl	Multivitamin VF	Francox	
	Vitamin AD3EK		